

# WIMPs and Other Particles Searches for

OMMITTED FROM SUMMARY TABLE  
A REVIEW GOES HERE – Check our WWW List of Reviews

## GALACTIC WIMP SEARCHES

### Cross-Section Limits for Dark Matter Particles ( $X^0$ ) on Nuclei

These limits are for weakly-interacting stable particles that may constitute the invisible mass in the galaxy. Unless otherwise noted, a local mass density of 0.3 GeV/cm<sup>3</sup> is assumed; see each paper for velocity distribution assumptions. In the papers the limit is given as a function of the  $X^0$  mass. Here we list limits only for typical mass values of 20 GeV, 100 GeV, and 1 TeV. Specific limits on supersymmetric dark matter particles may be found in the Supersymmetry section.

#### For $m_{X^0} = 20$ GeV

<u>VALUE (nb)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
1 AKERIB	06	CDMS	<sup>73</sup> Ge, <sup>29</sup> Si	
2 SHIMIZU	06A	CNTR	F (CaF <sub>2</sub> )	
3 ALNER	05	NAIA	NaI	
4 BARNABE-HE..05	PICA	F (C <sub>4</sub> F <sub>10</sub> )		
5 BENOIT	05	EDEL	<sup>73</sup> Ge	
6 GIRARD	05	SMPL	F (C <sub>2</sub> ClF <sub>5</sub> )	
7 Klapdor-K...05	HDMS	<sup>73</sup> Ge (enriched)		
8 MIUCHI	03	BOLO	LiF	
9 TAKEDA	03	BOLO	NaF	
< 0.08	90	10 ANGLOHER	02	CRES Al
		11 BENOIT	00	EDEL Ge
< 0.04	95	12 KLIMENTKO	98	CNTR <sup>73</sup> Ge, inel.
< 0.8		ALESSAND...	96	CNTR O
< 6		ALESSAND...	96	CNTR Te
< 0.02	90	13 BELLI	96	CNTR <sup>129</sup> Xe, inel.
		14 BELLI	96C	CNTR <sup>129</sup> Xe
< 0.004	90	15 BERNABEI	96	CNTR Na
< 0.3	90	15 BERNABEI	96	CNTR I
< 0.2	95	16 SARSA	96	CNTR Na
< 0.015	90	17 SMITH	96	CNTR Na
< 0.05	95	18 GARCIA	95	CNTR Natural Ge
< 0.1	95	QUENBY	95	CNTR Na
< 90	90	19 SNOWDEN...	95	MICA <sup>16</sup> O
< 4 × 10 <sup>3</sup>	90	19 SNOWDEN...	95	MICA <sup>39</sup> K
< 0.7	90	BACCI	92	CNTR Na
< 0.12	90	20 REUSSER	91	CNTR Natural Ge
< 0.06	95	CALDWELL	88	CNTR Natural Ge

- <sup>1</sup> AKERIB 06 give  $\sigma < 20$  (0.3) pb (90% CL) for spin-dependent  $X^0$ -proton (neutron) cross section. See also AKERIB 05.
- <sup>2</sup> SHIMIZU 06A give  $\sigma < 2$  (30) pb (90% CL) for spin-dependent  $X^0$ -proton (neutron) cross section.
- <sup>3</sup> ALNER 05 give  $\sigma < 0.5$  (60) pb (90% CL) for spin-dependent  $X^0$ -proton (neutron) cross section.
- <sup>4</sup> BARNABE-HEIDER 05 give  $\sigma < 1.5$  (20) pb (90% CL) for spin-dependent  $X^0$ -proton (neutron) cross section.
- <sup>5</sup> BENOIT 05 give  $\sigma < 10$  pb (90% CL) for spin-dependent  $X^0$ -neutron cross section.
- <sup>6</sup> GIRARD 05 give  $\sigma < 1.5$  pb (90% CL) for spin-dependent  $X^0$ -proton cross section.
- <sup>7</sup> Klapdor-Kleingrothaus 05 give  $\sigma < 4$  pb (90% CL) for spin-dependent  $X^0$ -neutron cross section.
- <sup>8</sup> MIUCHI 03 give model-independent limit  $\sigma < 35$  pb (90% CL) for spin-dependent  $X^0$ -proton cross section.
- <sup>9</sup> TAKEDA 03 give model-independent limit  $\sigma < 0.03$  (0.6) nb (90% CL) for spin-dependent  $X^0$ -proton (neutron) cross section.
- <sup>10</sup> ANGLOHER 02 limit is for spin-dependent WIMP-Aluminum cross section.
- <sup>11</sup> BENOIT 00 find four event categories in Ge detectors and suggest that low-energy surface nuclear recoils can explain anomalous events reported by UKDMC and Saclay NaI experiments.
- <sup>12</sup> KLIMENTKO 98 limit is for inelastic scattering  $X^0 \text{ } ^{73}\text{Ge} \rightarrow X^0 \text{ } ^{73}\text{Ge}^*$  (13.26 keV).
- <sup>13</sup> BELLI 96 limit for inelastic scattering  $X^0 \text{ } ^{129}\text{Xe} \rightarrow X^0 \text{ } ^{129}\text{Xe}^*$  (39.58 keV).
- <sup>14</sup> BELLI 96C use background subtraction and obtain  $\sigma < 150$  pb (< 1.5 fb) (90% CL) for spin-dependent (independent)  $X^0$ -proton cross section. The confidence level is from R. Bernabei, private communication, May 20, 1999.
- <sup>15</sup> BERNABEI 96 use pulse shape discrimination to enhance the possible signal. The limit here is from R. Bernabei, private communication, September 19, 1997.
- <sup>16</sup> SARSA 96 search for annual modulation of WIMP signal. See SARSA 97 for details of the analysis. The limit here is from M.L. Sarsa, private communication, May 26, 1997.
- <sup>17</sup> SMITH 96 use pulse shape discrimination to enhance the possible signal. A dark matter density of  $0.4 \text{ GeV cm}^{-3}$  is assumed.
- <sup>18</sup> GARCIA 95 limit is from the event rate. A weaker limit is obtained from searches for diurnal and annual modulation.
- <sup>19</sup> SNOWDEN-IFFT 95 look for recoil tracks in an ancient mica crystal. Similar limits are also given for  $^{27}\text{Al}$  and  $^{28}\text{Si}$ . See COLLAR 96 and SNOWDEN-IFFT 96 for discussion on potential backgrounds.
- <sup>20</sup> REUSSER 91 limit here is changed from published (0.04) after reanalysis by authors. J.L. Vuilleumier, private communication, March 29, 1996.

## For $m_{X^0} = 100 \text{ GeV}$

VALUE (nb)	CL%	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
21 AKERIB	06	CDMS	$^{73}\text{Ge}, ^{29}\text{Si}$	
22 SHIMIZU	06A	CNTR	F ( $\text{CaF}_2$ )	
23 ALNER	05	NAIA	NaI	
24 BARNABE-HE.05		PICA	F ( $\text{C}_4\text{F}_{10}$ )	
25 BENOIT	05	EDEL	$^{73}\text{Ge}$	
26 GIRARD	05	SMPL	F ( $\text{C}_2\text{ClF}_5$ )	
27 GIULIANI	05	RVUE		
28 GIULIANI	05A	RVUE		
29 Klapdor-K...05		HDMS	$^{73}\text{Ge}$ (enriched)	
30 GIULIANI	04	RVUE		
31 GIULIANI	04A	RVUE		

		32	MIUCHI	03	BOLO	LiF
		33	MIUCHI	03	BOLO	LiF
		34	TAKEDA	03	BOLO	NaF
< 0.3	90	35	ANGLOHER	02	CRES	AI
		36	BELLI	02	RVUE	
		37	BERNABEI	02C	DAMA	
		38	GREEN	02	RVUE	
		39	ULLIO	01	RVUE	
		40	BENOIT	00	EDEL	Ge
< 0.004	90	41	BERNABEI	00D		<sup>129</sup> Xe, inel.
		42	AMBROSIO	99	MCRO	
		43	BRHLIK	99	RVUE	
< 0.008	95	44	KLIMENKO	98	CNTR	<sup>73</sup> Ge, inel.
< 0.08	95	45	KLIMENKO	98	CNTR	<sup>73</sup> Ge, inel.
< 4			ALESSAND...	96	CNTR	O
< 25			ALESSAND...	96	CNTR	Te
< 0.006	90	46	BELLI	96	CNTR	<sup>129</sup> Xe, inel.
		47	BELLI	96C	CNTR	<sup>129</sup> Xe
< 0.001	90	48	BERNABEI	96	CNTR	Na
< 0.3	90	48	BERNABEI	96	CNTR	I
< 0.7	95	49	SARSA	96	CNTR	Na
< 0.03	90	50	SMITH	96	CNTR	Na
< 0.8	90	50	SMITH	96	CNTR	I
< 0.35	95	51	GARCIA	95	CNTR	Natural Ge
< 0.6	95		QUENBY	95	CNTR	Na
< 3	95		QUENBY	95	CNTR	I
< 1.5 × 10 <sup>2</sup>	90	52	SNOWDEN-...	95	MICA	<sup>16</sup> O
< 4 × 10 <sup>2</sup>	90	52	SNOWDEN-...	95	MICA	<sup>39</sup> K
< 0.08	90	53	BECK	94	CNTR	<sup>76</sup> Ge
< 2.5	90		BACCI	92	CNTR	Na
< 3	90		BACCI	92	CNTR	I
< 0.9	90	54	REUSSER	91	CNTR	Natural Ge
< 0.7	95		CALDWELL	88	CNTR	Natural Ge

<sup>21</sup> AKERIB 06 give  $\sigma < 5$  (0.07) pb (90% CL) for spin-dependent  $X^0$ -proton (neutron) cross section. See also AKERIB 05.

<sup>22</sup> SHIMIZU 06A give  $\sigma < 2$  (30) pb (90% CL) for spin-dependent  $X^0$ -proton (neutron) cross section.

<sup>23</sup> ALNER 05 give  $\sigma < 0.3$  (10) pb (90% CL) for spin-dependent  $X^0$ -proton (neutron) cross section.

<sup>24</sup> BARNABE-HEIDER 05 give  $\sigma < 2$  (30) pb (90% CL) for spin-dependent  $X^0$ -proton (neutron) cross section.

<sup>25</sup> BENOIT 05 give  $\sigma < 100$  (0.7) pb (90% CL) for spin-dependent  $X^0$ -proton (neutron) cross section.

<sup>26</sup> GIRARD 05 give  $\sigma < 1.5$  pb (90% CL) for spin-dependent  $X^0$ -proton cross section.

<sup>27</sup> GIULIANI 05 analyzes the spin-independent  $X^0$ -nucleon cross section limits with both isoscalar and isovector couplings. See Figs. 3 and 4 for limits on the couplings.

<sup>28</sup> GIULIANI 05A analyze available data and give combined limits  $\sigma < 0.7$  (0.2) pb for spin-dependent  $X^0$ -proton (neutron) cross section.

<sup>29</sup> Klapdor-Kleingrothaus 05 give  $\sigma < 1.5$  pb (90% CL) for spin-dependent  $X^0$ -neutron cross section.

<sup>30</sup> GIULIANI 04 reanalyze COLLAR 00 data and give limits for spin-dependent  $X^0$ -proton and neutron couplings.

- 31 GIULIANI 04A gives limits for spin-dependent  $X^0$ -proton and neutron couplings from existing data.
- 32 MIUCHI 03 give model-independent limit for spin-dependent  $X^0$ -proton and neutron cross sections. See their Fig. 5.
- 33 MIUCHI 03 give model-independent limit  $\sigma < 35 \text{ pb}$  (90% CL) for spin-dependent  $X^0$ -proton cross section.
- 34 TAKEDA 03 give model-independent limit  $\sigma < 0.04$  (0.8) nb (90% CL) for spin-dependent  $X^0$ -proton (neutron) cross section.
- 35 ANGLOHER 02 limit is for spin-dependent WIMP-Aluminum cross section.
- 36 BELLI 02 discuss dependence of the extracted WIMP cross section on the assumptions of the galactic halo structure.
- 37 BERNABEI 02C analyze the DAMA data in the scenario in which  $X^0$  scatters into a slightly heavier state as discussed by SMITH 01.
- 38 GREEN 02 discusses dependence of extracted WIMP cross section limits on the assumptions of the galactic halo structure.
- 39 ULLIO 01 disfavor the possibility that the BERNABEI 99 signal is due to spin-dependent WIMP coupling.
- 40 BENOIT 00 find four event categories in Ge detectors and suggest that low-energy surface nuclear recoils can explain anomalous events reported by UKDMC and Saclay NaI experiments.
- 41 BERNABEI 00D limit is for inelastic scattering  $X^0 \text{^{129}Xe} \rightarrow X^0 \text{^{129}Xe}$  (39.58 keV).
- 42 AMBROSIO 99 search for upgoing muon events induced by neutrinos originating from WIMP annihilations in the Sun and Earth.
- 43 BRHLIK 99 discuss the effect of astrophysical uncertainties on the WIMP interpretation of the BERNABEI 99 signal.
- 44 KLIMENKO 98 limit is for inelastic scattering  $X^0 \text{^{73}Ge} \rightarrow X^0 \text{^{73}Ge}^*$  (13.26 keV).
- 45 KLIMENKO 98 limit is for inelastic scattering  $X^0 \text{^{73}Ge} \rightarrow X^0 \text{^{73}Ge}^*$  (66.73 keV).
- 46 BELLI 96 limit for inelastic scattering  $X^0 \text{^{129}Xe} \rightarrow X^0 \text{^{129}Xe}^*$  (39.58 keV).
- 47 BELLI 96C use background subtraction and obtain  $\sigma < 0.35 \text{ pb}$  ( $< 0.15 \text{ fb}$ ) (90% CL) for spin-dependent (independent)  $X^0$ -proton cross section. The confidence level is from R. Bernabei, private communication, May 20, 1999.
- 48 BERNABEI 96 use pulse shape discrimination to enhance the possible signal. The limit here is from R. Bernabei, private communication, September 19, 1997.
- 49 SARSA 96 search for annual modulation of WIMP signal. See SARSA 97 for details of the analysis. The limit here is from M.L. Sarsa, private communication, May 26, 1997.
- 50 SMITH 96 use pulse shape discrimination to enhance the possible signal. A dark matter density of  $0.4 \text{ GeV cm}^{-3}$  is assumed.
- 51 GARCIA 95 limit is from the event rate. A weaker limit is obtained from searches for diurnal and annual modulation.
- 52 SNOWDEN-IFFT 95 look for recoil tracks in an ancient mica crystal. Similar limits are also given for  $^{27}\text{Al}$  and  $^{28}\text{Si}$ . See COLLAR 96 and SNOWDEN-IFFT 96 for discussion on potential backgrounds.
- 53 BECK 94 uses enriched  $^{76}\text{Ge}$  (86% purity).
- 54 REUSSER 91 limit here is changed from published (0.3) after reanalysis by authors. J.L. Vuilleumier, private communication, March 29, 1996.

**For  $m_{X^0} = 1 \text{ TeV}$** 

<u>VALUE (nb)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
55	AKERIB 06	CDMS	$^{73}\text{Ge}, ^{29}\text{Si}$	
56	ALNER 05	NAIA	Nal	
57	BARNABE-HE..05	PICA	F ( $\text{C}_4\text{F}_{10}$ )	
58	BENOIT 05	EDEL	$^{73}\text{Ge}$	
59	GIRARD 05	SMPL	F ( $\text{C}_2\text{ClF}_5$ )	
60	KLAPDOR-K...05	HDMs	$^{73}\text{Ge}$ (enriched)	
61	MIUCHI 03	BOLO	LiF	
62	TAKEDA 03	BOLO	NaF	
< 3	90	ANGLOHER 02	CRES	Al
		64 BENOIT 00	EDEL	Ge
		65 BERNABEI 99D	CNTR	SIMP
		66 DERBIN 99	CNTR	SIMP
< 0.06	95	67 KLIMENKO 98	CNTR	$^{73}\text{Ge}$ , inel.
< 0.4	95	68 KLIMENKO 98	CNTR	$^{73}\text{Ge}$ , inel.
< 40		ALESSAND... 96	CNTR	O
<700		ALESSAND... 96	CNTR	Te
< 0.05	90	69 BELLI 96	CNTR	$^{129}\text{Xe}$ , inel.
< 1.5	90	70 BELLI 96	CNTR	$^{129}\text{Xe}$ , inel.
		71 BELLI 96C	CNTR	$^{129}\text{Xe}$
< 0.01	90	72 BERNABEI 96	CNTR	Na
< 9	90	72 BERNABEI 96	CNTR	I
< 7	95	73 SARSA 96	CNTR	Na
< 0.3	90	74 SMITH 96	CNTR	Na
< 6	90	74 SMITH 96	CNTR	I
< 6	95	75 GARCIA 95	CNTR	Natural Ge
< 8	95	QUENBY 95	CNTR	Na
< 50	95	QUENBY 95	CNTR	I
< $7 \times 10^2$	90	76 SNOWDEN-... 95	MICA	$^{16}\text{O}$
< $1 \times 10^3$	90	76 SNOWDEN-... 95	MICA	$^{39}\text{K}$
< 0.8	90	77 BECK 94	CNTR	$^{76}\text{Ge}$
< 30	90	BACCI 92	CNTR	Na
< 30	90	BACCI 92	CNTR	I
< 15	90	78 REUSSER 91	CNTR	Natural Ge
< 6	95	CALDWELL 88	CNTR	Natural Ge

55 AKERIB 06 give  $\sigma < 30$  (0.5) pb (90% CL) for spin-dependent  $X^0$ -proton (neutron) cross section. See also AKERIB 05.

56 ALNER 05 give  $\sigma < 1.5$  (40) pb (90% CL) for spin-dependent  $X^0$ -proton (neutron) cross section.

57 BARNABE-HEIDER 05 give  $\sigma < 15$  (200) pb (90% CL) for spin-dependent  $X^0$ -proton (neutron) cross section.

58 BENOIT 05 give  $\sigma < 600$  (4) pb (90% CL) for spin-dependent  $X^0$ -proton (neutron) cross section.

59 GIRARD 05 give  $\sigma < 10$  pb (90% CL) for spin-dependent  $X^0$ -proton cross section.

60 Klapdor-Kleingrothaus 05 give  $\sigma < 10$  pb (90% CL) for spin-dependent  $X^0$ -neutron cross section.

61 MIUCHI 03 give model-independent limit  $\sigma < 260$  pb (90% CL) for spin-dependent  $X^0$ -proton cross section.

- 62 TAKEDA 03 give model-independent limit  $\sigma < 0.15$  (4) nb (90% CL) for spin-dependent  $X^0$ -proton (neutron) cross section.
- 63 ANGLOHER 02 limit is for spin-dependent WIMP-Aluminum cross section.
- 64 BENOIT 00 find four event categories in Ge detectors and suggest that low-energy surface nuclear recoils can explain anomalous events reported by UKDMC and Saclay NaI experiments.
- 65 BERNABEI 99D search for SIMPs (Strongly Interacting Massive Particles) in the mass range  $10^3$ – $10^{16}$  GeV. See their Fig. 3 for cross-section limits.
- 66 DERBIN 99 search for SIMPs (Strongly Interacting Massive Particles) in the mass range  $10^2$ – $10^{14}$  GeV. See their Fig. 3 for cross-section limits.
- 67 KLIMENKO 98 limit is for inelastic scattering  $X^0$   $^{73}\text{Ge} \rightarrow X^0$   $^{73}\text{Ge}^*$  (13.26 keV).
- 68 KLIMENKO 98 limit is for inelastic scattering  $X^0$   $^{73}\text{Ge} \rightarrow X^0$   $^{73}\text{Ge}^*$  (66.73 keV).
- 69 BELLI 96 limit for inelastic scattering  $X^0$   $^{129}\text{Xe} \rightarrow X^0$   $^{129}\text{Xe}^*$  (39.58 keV).
- 70 BELLI 96 limit for inelastic scattering  $X^0$   $^{129}\text{Xe} \rightarrow X^0$   $^{129}\text{Xe}^*$  (236.14 keV).
- 71 BELLI 96C use background subtraction and obtain  $\sigma < 0.7$  pb (< 0.7 fb) (90% CL) for spin-dependent (independent)  $X^0$ -proton cross section. The confidence level is from R. Bernabei, private communication, May 20, 1999.
- 72 BERNABEI 96 use pulse shape discrimination to enhance the possible signal. The limit here is from R. Bernabei, private communication, September 19, 1997.
- 73 SARSA 96 search for annual modulation of WIMP signal. See SARSA 97 for details of the analysis. The limit here is from M.L. Sarsa, private communication, May 26, 1997.
- 74 SMITH 96 use pulse shape discrimination to enhance the possible signal. A dark matter density of  $0.4 \text{ GeV cm}^{-3}$  is assumed.
- 75 GARCIA 95 limit is from the event rate. A weaker limit is obtained from searches for diurnal and annual modulation.
- 76 SNOWDEN-IFFT 95 look for recoil tracks in an ancient mica crystal. Similar limits are also given for  $^{27}\text{Al}$  and  $^{28}\text{Si}$ . See COLLAR 96 and SNOWDEN-IFFT 96 for discussion on potential backgrounds.
- 77 BECK 94 uses enriched  $^{76}\text{Ge}$  (86% purity).
- 78 REUSSER 91 limit here is changed from published (5) after reanalysis by authors. J.L. Vuilleumier, private communication, March 29, 1996.

## CONCENTRATION OF STABLE PARTICLES IN MATTER

### Concentration of Heavy (Charge +1) Stable Particles in Matter

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
$<4 \times 10^{-17}$	95	79 YAMAGATA	93	SPEC Deep sea water, $M=5\text{--}1600m_p$
$<6 \times 10^{-15}$	95	80 VERKERK	92	SPEC Water, $M=10^5$ to $3 \times 10^7$ GeV
$<7 \times 10^{-15}$	95	80 VERKERK	92	SPEC Water, $M=10^4$ , $6 \times 10^7$ GeV
$<9 \times 10^{-15}$	95	80 VERKERK	92	SPEC Water, $M=10^8$ GeV
$<3 \times 10^{-23}$	90	81 HEMMICK	90	SPEC Water, $M=1000m_p$
$<2 \times 10^{-21}$	90	81 HEMMICK	90	SPEC Water, $M=5000m_p$
$<3 \times 10^{-20}$	90	81 HEMMICK	90	SPEC Water, $M=10000m_p$
$<1. \times 10^{-29}$		SMITH	82B	SPEC Water, $M=30\text{--}400m_p$
$<2. \times 10^{-28}$		SMITH	82B	SPEC Water, $M=12\text{--}1000m_p$
$<1. \times 10^{-14}$		SMITH	82B	SPEC Water, $M>1000 m_p$
$<(0.2\text{--}1.) \times 10^{-21}$		SMITH	79	SPEC Water, $M=6\text{--}350 m_p$

79 YAMAGATA 93 used deep sea water at 4000 m since the concentration is enhanced in deep sea due to gravity.

80 VERKERK 92 looked for heavy isotopes in sea water and put a bound on concentration of stable charged massive particle in sea water. The above bound can be translated into a bound on charged dark matter particle ( $5 \times 10^6$  GeV), assuming the local density,  $\rho=0.3$  GeV/cm<sup>3</sup>, and the mean velocity  $\langle v \rangle=300$  km/s.

81 See HEMMICK 90 Fig. 7 for other masses 100–10000  $m_p$ .

## Concentration of Heavy Stable Particles Bound to Nuclei

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
<1.2 × 10 <sup>-11</sup>	95	82 JAVORSEK	01	SPEC Au, $M= 3$ GeV
<6.9 × 10 <sup>-10</sup>	95	82 JAVORSEK	01	SPEC Au, $M= 144$ GeV
<1 × 10 <sup>-11</sup>	95	83 JAVORSEK	01B	SPEC Au, $M= 188$ GeV
<1 × 10 <sup>-8</sup>	95	83 JAVORSEK	01B	SPEC Au, $M= 1669$ GeV
<6 × 10 <sup>-9</sup>	95	83 JAVORSEK	01B	SPEC Fe, $M= 188$ GeV
<1 × 10 <sup>-8</sup>	95	83 JAVORSEK	01B	SPEC Fe, $M= 647$ GeV
<4 × 10 <sup>-20</sup>	90	84 HEMMICK	90	SPEC C, $M= 100m_p$
<8 × 10 <sup>-20</sup>	90	84 HEMMICK	90	SPEC C, $M= 1000m_p$
<2 × 10 <sup>-16</sup>	90	84 HEMMICK	90	SPEC C, $M= 10000m_p$
<6 × 10 <sup>-13</sup>	90	84 HEMMICK	90	SPEC Li, $M= 1000m_p$
<1 × 10 <sup>-11</sup>	90	84 HEMMICK	90	SPEC Be, $M= 1000m_p$
<6 × 10 <sup>-14</sup>	90	84 HEMMICK	90	SPEC B, $M= 1000m_p$
<4 × 10 <sup>-17</sup>	90	84 HEMMICK	90	SPEC O, $M= 1000m_p$
<4 × 10 <sup>-15</sup>	90	84 HEMMICK	90	SPEC F, $M= 1000m_p$
<1.5 × 10 <sup>-13</sup> /nucleon	68	85 NORMAN	89	SPEC $^{206}\text{Pb}X^-$
<1.2 × 10 <sup>-12</sup> /nucleon	68	85 NORMAN	87	SPEC $^{56,58}\text{Fe}X^-$

82 JAVORSEK 01 search for (neutral) SIMPs (strongly interacting massive particles) bound to Au nuclei. Here  $M$  is the effective SIMP mass.

83 JAVORSEK 01B search for (neutral) SIMPs (strongly interacting massive particles) bound to Au and Fe nuclei from various origins with exposures on the earth's surface, in a satellite, heavy ion collisions, etc. Here  $M$  is the mass of the anomalous nucleus. See also JAVORSEK 02.

84 See HEMMICK 90 Fig. 7 for other masses 100–10000  $m_p$ .

85 Bound valid up to  $m_{X^-} \sim 100$  TeV.

## LIMITS ON NEUTRAL PARTICLE PRODUCTION

### Production Cross Section of Radiatively-Decaying Neutral Particle

VALUE (pb)	CL%	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
<(0.043–0.17)	95	86 ABBIENDI	00D OPAL	$e^+e^- \rightarrow X^0Y^0,$ $X^0 \rightarrow Y^0\gamma$
<(0.05–0.8)	95	87 ABBIENDI	00D OPAL	$e^+e^- \rightarrow X^0X^0,$ $X^0 \rightarrow Y^0\gamma$
<(2.5–0.5)	95	88 ACKERSTAFF	97B OPAL	$e^+e^- \rightarrow X^0Y^0,$ $X^0 \rightarrow Y^0\gamma$
<(1.6–0.9)	95	89 ACKERSTAFF	97B OPAL	$e^+e^- \rightarrow X^0X^0,$ $X^0 \rightarrow Y^0\gamma$

- 86 ABBIENDI 00D associated production limit is for  $m_{X^0} = 90\text{--}188$  GeV,  $m_{Y^0}=0$  at  $E_{cm}=189$  GeV. See also their Fig. 9.
- 87 ABBIENDI 00D pair production limit is for  $m_{X^0} = 45\text{--}94$  GeV,  $m_{Y^0}=0$  at  $E_{cm}=189$  GeV. See also their Fig. 12.
- 88 ACKERSTAFF 97B associated production limit is for  $m_{X^0} = 80\text{--}160$  GeV,  $m_{Y^0}=0$  from  $10.0 \text{ pb}^{-1}$  at  $E_{cm} = 161$  GeV. See their Fig. 3(a).
- 89 ACKERSTAFF 97B pair production limit is for  $m_{X^0} = 40\text{--}80$  GeV,  $m_{Y^0}=0$  from  $10.0 \text{ pb}^{-1}$  at  $E_{cm} = 161$  GeV. See their Fig. 3(b).

## Heavy Particle Production Cross Section

<u>VALUE (cm<sup>2</sup>/N)</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>					
			90 ADAMS	97B KTEV	$m = 1.2\text{--}5$ GeV
$< 10^{-36}\text{--}10^{-33}$	90		91 GALLAS	95 TOF	$m = 0.5\text{--}20$ GeV
$<(4\text{--}0.3) \times 10^{-31}$	95		92 AKESSON	91 CNTR	$m = 0\text{--}5$ GeV
$<2 \times 10^{-36}$	90	0	93 BADIER	86 BDMP	$\tau = (0.05\text{--}1.) \times 10^{-8}$ s
$<2.5 \times 10^{-35}$		0	94 GUSTAFSON	76 CNTR	$\tau > 10^{-7}$ s

- 90 ADAMS 97B search for a hadron-like neutral particle produced in  $pN$  interactions, which decays into a  $\rho^0$  and a weakly interacting massive particle. Upper limits are given for the ratio to  $K_L$  production for the mass range 1.2–5 GeV and lifetime  $10^{-9}\text{--}10^{-4}$  s. See also our Light Gluino Section.
- 91 GALLAS 95 limit is for a weakly interacting neutral particle produced in 800 GeV/c  $pN$  interactions decaying with a lifetime of  $10^{-4}\text{--}10^{-8}$  s. See their Figs. 8 and 9. Similar limits are obtained for a stable particle with interaction cross section  $10^{-29}\text{--}10^{-33}$  cm<sup>2</sup>. See Fig. 10.
- 92 AKESSON 91 limit is from weakly interacting neutral long-lived particles produced in  $pN$  reaction at 450 GeV/c performed at CERN SPS. Bourquin-Gaillard formula is used as the production model. The above limit is for  $\tau > 10^{-7}$  s. For  $\tau > 10^{-9}$  s,  $\sigma < 10^{-30}$  cm<sup>-2</sup>/nucleon is obtained.
- 93 BADIER 86 looked for long-lived particles at 300 GeV  $\pi^-$  beam dump. The limit applies for nonstrongly interacting neutral or charged particles with mass  $>2$  GeV. The limit applies for particle modes,  $\mu^+ \pi^-$ ,  $\mu^+ \mu^-$ ,  $\pi^+ \pi^- X$ ,  $\pi^+ \pi^- \pi^\pm$  etc. See their figure 5 for the contours of limits in the mass- $\tau$  plane for each mode.
- 94 GUSTAFSON 76 is a 300 GeV FNAL experiment looking for heavy ( $m > 2$  GeV) long-lived neutral hadrons in the M4 neutral beam. The above typical value is for  $m = 3$  GeV and assumes an interaction cross section of 1 mb. Values as a function of mass and interaction cross section are given in figure 2.

## Production of New Penetrating Non- $\nu$ Like States in Beam Dump

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
95 LOSECCO	81 CALO	28 GeV protons	
95 No excess neutral-current events leads to $\sigma(\text{production}) \times \sigma(\text{interaction}) \times \text{acceptance} < 2.26 \times 10^{-71}$ cm <sup>4</sup> /nucleon <sup>2</sup> (CL = 90%) for light neutrals. Acceptance depends on models (0.1 to $4 \times 10^{-4}$ ).			

## LIMITS ON JET-JET RESONANCES

### Heavy Particle Production Cross Section in $p\bar{p}$

Limits are for a particle decaying to two hadronic jets.

<u>Units(pb)</u>	<u>CL%</u>	<u>Mass(GeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>					
			96 ABE	99F CDF	1.8 TeV $p\bar{p} \rightarrow b\bar{b}$ + anything
			97 ABE	97G CDF	1.8 TeV $p\bar{p} \rightarrow 2$ jets
<2603	95	200	98 ABE	93G CDF	1.8 TeV $p\bar{p} \rightarrow 2$ jets
< 44	95	400	98 ABE	93G CDF	1.8 TeV $p\bar{p} \rightarrow 2$ jets
< 7	95	600	98 ABE	93G CDF	1.8 TeV $p\bar{p} \rightarrow 2$ jets
96 ABE 99F search for narrow $b\bar{b}$ resonances in $p\bar{p}$ collisions at $E_{cm}=1.8$ TeV. Limits on $\sigma(p\bar{p} \rightarrow X + \text{anything}) \times B(X \rightarrow b\bar{b})$ in the range $3-10^3$ pb (95%CL) are given for $m_X=200-750$ GeV. See their Table I.					
97 ABE 97G search for narrow dijet resonances in $p\bar{p}$ collisions with $106 \text{ pb}^{-1}$ of data at $E_{cm} = 1.8$ TeV. Limits on $\sigma(p\bar{p} \rightarrow X + \text{anything}) \cdot B(X \rightarrow jj)$ in the range $10^4-10^{-1}$ pb (95%CL) are given for dijet mass $m=200-1150$ GeV with both jets having $ \eta  < 2.0$ and the dijet system having $ \cos\theta^*  < 0.67$ . See their Table I for the list of limits. Supersedes ABE 93G.					
98 ABE 93G gives cross section times branching ratio into light ( $d, u, s, c, b$ ) quarks for $\Gamma = 0.02 M$ . Their Table II gives limits for $M = 200-900$ GeV and $\Gamma = (0.02-0.2) M$ .					

## LIMITS ON CHARGED PARTICLES IN $e^+ e^-$

### Heavy Particle Production Cross Section in $e^+ e^-$

Ratio to  $\sigma(e^+ e^- \rightarrow \mu^+ \mu^-)$  unless noted. See also entries in Free Quark Search and Magnetic Monopole Searches.

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>					
			99 ACKERSTAFF	98P OPAL	$Q=1,2/3, m=45-89.5$ GeV
			100 ABREU	97D DLPH	$Q=1,2/3, m=45-84$ GeV
			101 BARATE	97K ALEP	$Q=1, m=45-85$ GeV
$<2 \times 10^{-5}$	95		102 AKERS	95R OPAL	$Q=1, m=5-45$ GeV
$<1 \times 10^{-5}$	95		102 AKERS	95R OPAL	$Q=2, m=5-45$ GeV
$<2 \times 10^{-3}$	90		103 BUSKULIC	93C ALEP	$Q=1, m=32-72$ GeV
$<(10^{-2}-1)$	95		104 ADACHI	90C TOPZ	$Q = 1, m = 1-16, 18-27$ GeV
$<7 \times 10^{-2}$	90		105 ADACHI	90E TOPZ	$Q = 1, m = 5-25$ GeV
$<1.6 \times 10^{-2}$	95	0	106 KINOSHITA	82 PLAS	$Q=3-180, m < 14.5$ GeV
$<5.0 \times 10^{-2}$	90	0	107 BARTEL	80 JADE	$Q=(3,4,5)/3$ 2-12 GeV
99 ACKERSTAFF 98P search for pair production of long-lived charged particles at $E_{cm}$ between 130 and 183 GeV and give limits $\sigma < (0.05-0.2)$ pb (95%CL) for spin-0 and spin-1/2 particles with $m=45-89.5$ GeV, charge 1 and 2/3. The limit is translated to the cross section at $E_{cm}=183$ GeV with the $s$ dependence described in the paper. See their Figs. 2-4.					
100 ABREU 97D search for pair production of long-lived particles and give limits $\sigma < (0.4-2.3)$ pb (95%CL) for various center-of-mass energies $E_{cm}=130-136, 161$ , and 172 GeV, assuming an almost flat production distribution in $\cos\theta$ .					

- 101 BARATE 97K search for pair production of long-lived charged particles at  $E_{\text{cm}} = 130, 136, 161,$  and  $172 \text{ GeV}$  and give limits  $\sigma < (0.2\text{--}0.4) \text{ pb}$  (95%CL) for spin-0 and spin-1/2 particles with  $m=45\text{--}85 \text{ GeV}$ . The limit is translated to the cross section at  $E_{\text{cm}}=172 \text{ GeV}$  with the  $E_{\text{cm}}$  dependence described in the paper. See their Figs. 2 and 3 for limits on  $J = 1/2$  and  $J = 0$  cases.
- 102 AKERS 95R is a CERN-LEP experiment with  $W_{\text{cm}} \sim m_Z$ . The limit is for the production of a stable particle in multihadron events normalized to  $\sigma(e^+e^- \rightarrow \text{hadrons})$ . Constant phase space distribution is assumed. See their Fig. 3 for bounds for  $Q = \pm 2/3, \pm 4/3$ .
- 103 BUSKULIC 93C is a CERN-LEP experiment with  $W_{\text{cm}} = m_Z$ . The limit is for a pair or single production of heavy particles with unusual ionization loss in TPC. See their Fig. 5 and Table 1.
- 104 ADACHI 90C is a KEK-TRISTAN experiment with  $W_{\text{cm}} = 52\text{--}60 \text{ GeV}$ . The limit is for pair production of a scalar or spin-1/2 particle. See Figs. 3 and 4.
- 105 ADACHI 90E is KEK-TRISTAN experiment with  $W_{\text{cm}} = 52\text{--}61.4 \text{ GeV}$ . The above limit is for inclusive production cross section normalized to  $\sigma(e^+e^- \rightarrow \mu^+\mu^-)\cdot\beta(3-\beta^2)/2$ , where  $\beta = (1 - 4m^2/W_{\text{cm}}^2)^{1/2}$ . See the paper for the assumption about the production mechanism.
- 106 KINOSHITA 82 is SLAC PEP experiment at  $W_{\text{cm}} = 29 \text{ GeV}$  using lexan and  $^{39}\text{Cr}$  plastic sheets sensitive to highly ionizing particles.
- 107 BARTEL 80 is DESY-PETRA experiment with  $W_{\text{cm}} = 27\text{--}35 \text{ GeV}$ . Above limit is for inclusive pair production and ranges between  $1. \times 10^{-1}$  and  $1. \times 10^{-2}$  depending on mass and production momentum distributions. (See their figures 9, 10, 11).

### Branching Fraction of $Z^0$ to a Pair of Stable Charged Heavy Fermions

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
$<5 \times 10^{-6}$	95	108 AKERS	95R OPAL	$m = 40.4\text{--}45.6 \text{ GeV}$
$<1 \times 10^{-3}$	95	AKRAWY	900 OPAL	$m = 29\text{--}40 \text{ GeV}$
<b>108 AKERS 95R give the 95% CL limit <math>\sigma(X\bar{X})/\sigma(\mu\mu) &lt; 1.8 \times 10^{-4}</math> for the pair production of singly- or doubly-charged stable particles. The limit applies for the mass range 40.4–45.6 GeV for <math>X^\pm</math> and <math>&lt; 45.6 \text{ GeV}</math> for <math>X^{\pm\pm}</math>. See the paper for bounds for <math>Q = \pm 2/3, \pm 4/3</math>.</b>				

## LIMITS ON CHARGED PARTICLES IN HADRONIC REACTIONS

### Heavy Particle Production Cross Section

VALUE (nb)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>					
$<0.19$	95	109 AKTAS	04C H1	$m=3\text{--}10 \text{ GeV}$	
$<0.05$	95	110 ABE	92J CDF	$m=50\text{--}200 \text{ GeV}$	
$<30\text{--}130$		111 CARROLL	78 SPEC	$m=2\text{--}2.5 \text{ GeV}$	
$<100$	0	112 LEIPUNER	73 CNTR	$m=3\text{--}11 \text{ GeV}$	
<b>109 AKTAS 04C look for charged particle photoproduction at HERA with mean c.m. energy of 200 GeV.</b>					
<b>110 ABE 92J look for pair production of unit-charged particles which leave detector before decaying. Limit shown here is for <math>m=50 \text{ GeV}</math>. See their Fig. 5 for different charges and stronger limits for higher mass.</b>					
<b>111 CARROLL 78 look for neutral, <math>S = -2</math> dihyperon resonance in <math>p p \rightarrow 2K^+ X</math>. Cross section varies within above limits over mass range and <math>p_{\text{lab}} = 5.1\text{--}5.9 \text{ GeV}/c</math>.</b>					
<b>112 LEIPUNER 73 is an NAL 300 GeV <math>p</math> experiment. Would have detected particles with lifetime greater than 200 ns.</b>					

## Heavy Particle Production Differential Cross Section

VALUE (cm <sup>2</sup> sr <sup>-1</sup> GeV <sup>-1</sup> )	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>						
<2.6 × 10 <sup>-36</sup>	90	0	113 BALDIN	76	CNTR	—
						Q= 1, $m=2.1\text{--}9.4$ GeV
<2.2 × 10 <sup>-33</sup>	90	0	114 ALBROW	75	SPEC	±
						Q= ±1, $m=4\text{--}15$ GeV
<1.1 × 10 <sup>-33</sup>	90	0	114 ALBROW	75	SPEC	±
						Q= ±2, $m=6\text{--}27$ GeV
<8. × 10 <sup>-35</sup>	90	0	115 JOVANOV...	75	CNTR	±
						$m=15\text{--}26$ GeV
<1.5 × 10 <sup>-34</sup>	90	0	115 JOVANOV...	75	CNTR	±
						Q= ±2, $m=3\text{--}10$ GeV
<6. × 10 <sup>-35</sup>	90	0	115 JOVANOV...	75	CNTR	±
						Q= ±2, $m=10\text{--}26$ GeV
<1. × 10 <sup>-31</sup>	90	0	116 APPEL	74	CNTR	±
						$m=3.2\text{--}7.2$ GeV
<5.8 × 10 <sup>-34</sup>	90	0	117 ALPER	73	SPEC	±
						$m=1.5\text{--}24$ GeV
<1.2 × 10 <sup>-35</sup>	90	0	118 ANTIPOV	71B	CNTR	—
						Q= —, $m=2.2\text{--}2.8$
<2.4 × 10 <sup>-35</sup>	90	0	119 ANTIPOV	71C	CNTR	—
						Q= —, $m=1.2\text{--}1.7$ , 2.1–4
<2.4 × 10 <sup>-35</sup>	90	0	BINON	69	CNTR	—
						Q= —, $m=1\text{--}1.8$ GeV
<1.5 × 10 <sup>-36</sup>		0	120 DORFAN	65	CNTR	
						Be target $m=3\text{--}7$ GeV
<3.0 × 10 <sup>-36</sup>		0	120 DORFAN	65	CNTR	
						Fe target $m=3\text{--}7$ GeV

113 BALDIN 76 is a 70 GeV Serpukhov experiment. Value is per Al nucleus at  $\theta = 0$ . For other charges in range –0.5 to –3.0, CL = 90% limit is  $(2.6 \times 10^{-36}) / |(\text{charge})|$  for mass range (2.1–9.4 GeV)  $\times |(\text{charge})|$ . Assumes stable particle interacting with matter as do antiprotons.

114 ALBROW 75 is a CERN ISR experiment with  $E_{\text{cm}} = 53$  GeV.  $\theta = 40$  mr. See figure 5 for mass ranges up to 35 GeV.

115 JOVANOVICH 75 is a CERN ISR 26+26 and 15+15 GeV  $p p$  experiment. Figure 4 covers ranges  $Q = 1/3$  to 2 and  $m = 3$  to 26 GeV. Value is per GeV momentum.

116 APPEL 74 is NAL 300 GeV  $pW$  experiment. Studies forward production of heavy (up to 24 GeV) charged particles with momenta 24–200 GeV (–charge) and 40–150 GeV (+charge). Above typical value is for 75 GeV and is per GeV momentum per nucleon.

117 ALPER 73 is CERN ISR 26+26 GeV  $p p$  experiment.  $p > 0.9$  GeV,  $0.2 < \beta < 0.65$ .

118 ANTIPOV 71B is from same 70 GeV  $p$  experiment as ANTIPOV 71C and BINON 69.

119 ANTIPOV 71C limit inferred from flux ratio. 70 GeV  $p$  experiment.

120 DORFAN 65 is a 30 GeV/c  $p$  experiment at BNL. Units are per GeV momentum per nucleus.

## Long-Lived Heavy Particle Invariant Cross Section

VALUE (cm <sup>2</sup> /GeV <sup>2</sup> /N)	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>					
<5–700 × 10 <sup>-35</sup>	90	121 BERNSTEIN	88	CNTR	
<5–700 × 10 <sup>-37</sup>	90	121 BERNSTEIN	88	CNTR	
<2.5 × 10 <sup>-36</sup>	90	122 THRON	85	CNTR	—
<1. × 10 <sup>-35</sup>	90	122 THRON	85	CNTR	+
<6. × 10 <sup>-33</sup>	90	123 ARMITAGE	79	SPEC	$Q= 1, m=4\text{--}12$ GeV
					$m=1.87$ GeV

$<1.5 \times 10^{-33}$	90	123 ARMITAGE 124 BOZZOLI	79	SPEC CNTR $\pm$	$m=1.5\text{--}3.0 \text{ GeV}$ $Q = (2/3, 1, 4/3, 2)$
$<1.1 \times 10^{-37}$	90	125 CUTTS	78	CNTR	$m=4\text{--}10 \text{ GeV}$
$<3.0 \times 10^{-37}$	90	126 VIDAL	78	CNTR	$m=4.5\text{--}6 \text{ GeV}$

121 BERNSTEIN 88 limits apply at  $x = 0.2$  and  $p_T = 0$ . Mass and lifetime dependence of limits are shown in the regions:  $m = 1.5\text{--}7.5 \text{ GeV}$  and  $\tau = 10^{-8}\text{--}2 \times 10^{-6} \text{ s}$ . First number is for hadrons; second is for weakly interacting particles.

122 THRON 85 is FNAL 400 GeV proton experiment. Mass determined from measured velocity and momentum. Limits are for  $\tau > 3 \times 10^{-9} \text{ s}$ .

123 ARMITAGE 79 is CERN-ISR experiment at  $E_{\text{cm}} = 53 \text{ GeV}$ . Value is for  $x = 0.1$  and  $p_T = 0.15$ . Observed particles at  $m = 1.87 \text{ GeV}$  are found all consistent with being antideuterons.

124 BOZZOLI 79 is CERN-SPS 200 GeV  $pN$  experiment. Looks for particle with  $\tau$  larger than  $10^{-8} \text{ s}$ . See their figure 11–18 for production cross-section upper limits vs mass.

125 CUTTS 78 is  $pBe$  experiment at FNAL sensitive to particles of  $\tau > 5 \times 10^{-8} \text{ s}$ . Value is for  $-0.3 < x < 0$  and  $p_T = 0.175$ .

126 VIDAL 78 is FNAL 400 GeV proton experiment. Value is for  $x = 0$  and  $p_T = 0$ . Puts lifetime limit of  $< 5 \times 10^{-8} \text{ s}$  on particle in this mass range.

## Long-Lived Heavy Particle Production ( $\sigma(\text{Heavy Particle}) / \sigma(\pi)$ )

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$<10^{-8}$		127 NAKAMURA 0 128 BUSSIERE	89 80	SPEC $\pm$ CNTR $\pm$	$Q = (-5/3, \pm 2)$ $Q = (2/3, 1, 4/3, 2)$

127 NAKAMURA 89 is KEK experiment with 12 GeV protons on Pt target. The limit applies for mass  $\lesssim 1.6 \text{ GeV}$  and lifetime  $\gtrsim 10^{-7} \text{ s}$ .

128 BUSSIERE 80 is CERN-SPS experiment with 200–240 GeV protons on Be and Al target. See their figures 6 and 7 for cross-section ratio vs mass.

## Production and Capture of Long-Lived Massive Particles

VALUE ( $10^{-36} \text{ cm}^2$ )	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$				
$<20$ to $800$	0	129 ALEKSEEV	76	ELEC $\tau = 5 \text{ ms to 1 day}$
$<200$ to $2000$	0	129 ALEKSEEV	76B	ELEC $\tau = 100 \text{ ms to 1 day}$
$<1.4$ to $9$	0	130 FRANKEL	75	CNTR $\tau = 50 \text{ ms to 10 hours}$
$<0.1$ to $9$	0	131 FRANKEL	74	CNTR $\tau = 1 \text{ to 1000 hours}$

129 ALEKSEEV 76 and ALEKSEEV 76B are 61–70 GeV  $p$  Serpukhov experiment. Cross section is per Pb nucleus.

130 FRANKEL 75 is extension of FRANKEL 74.

131 FRANKEL 74 looks for particles produced in thick Al targets by 300–400 GeV/c protons.

## Long-Lived Particle Search at Hadron Collisions

Limits are for cross section times branching ratio.

VALUE (pb/nucleon)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$\bullet \bullet \bullet$ We do not use the following data for averages, fits, limits, etc. $\bullet \bullet \bullet$					
$<2$	90	0	132 BADIER	86	BDMP $\tau = (0.05\text{--}1.) \times 10^{-8} \text{ s}$
132 BADIER 86 looked for long-lived particles at 300 GeV $\pi^-$ beam dump. The limit applies for nonstrongly interacting neutral or charged particles with mass $> 2 \text{ GeV}$ . The limit applies for particle modes, $\mu^+ \pi^-$ , $\mu^+ \mu^-$ , $\pi^+ \pi^- X$ , $\pi^+ \pi^- \pi^\pm$ etc. See their figure 5 for the contours of limits in the mass- $\tau$ plane for each mode.					

## Long-Lived Heavy Particle Cross Section

<u>VALUE</u> (pb/sr)	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
<34	95	133 RAM	94	SPEC $1015 < m_{X^{++}} < 1085$ MeV
<75	95	133 RAM	94	SPEC $920 < m_{X^{++}} < 1025$ MeV
133 RAM 94 search for a long-lived doubly-charged fermion $X^{++}$ with mass between $m_N$ and $m_N + m_\pi$ and baryon number +1 in the reaction $pp \rightarrow X^{++} n$ . No candidate is found. The limit is for the cross section at $15^\circ$ scattering angle at 460 MeV incident energy and applies for $\tau(X^{++}) \gg 0.1 \mu s$ .				

## LIMITS ON CHARGED PARTICLES IN COSMIC RAYS

### Heavy Particle Flux in Cosmic Rays

<u>VALUE</u> ( $\text{cm}^{-2} \text{sr}^{-1} \text{s}^{-1}$ )	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>						
$\sim 6 \times 10^{-9}$		2	134 SAITO	90		$Q \simeq 14, m \simeq 370 m_p$
$< 1.4 \times 10^{-12}$	90	0	135 MINCER 136 SAKUYAMA	85 83B	CALO PLAS	$m \geq 1 \text{ TeV}$ $m \sim 1 \text{ TeV}$
$< 1.7 \times 10^{-11}$	99	0	137 BHAT	82	CC	
$< 1. \times 10^{-9}$	90	0	138 MARINI	82	CNTR $\pm$	$Q=1, m \sim 4.5 m_p$
2. $\times 10^{-9}$		3	139 YOCK	81	SPRK $\pm$	$Q=1, m \sim 4.5 m_p$
		3	139 YOCK	81	SPRK	Fractionally charged
3.0 $\times 10^{-9}$		3	140 YOCK	80	SPRK	$m \sim 4.5 m_p$
$(4 \pm 1) \times 10^{-11}$		3	GOODMAN	79	ELEC	$m \geq 5 \text{ GeV}$
$< 1.3 \times 10^{-9}$	90		141 BHAT	78	CNTR $\pm$	$m > 1 \text{ GeV}$
$< 1.0 \times 10^{-9}$		0	BRIATORE	76	ELEC	
$< 7. \times 10^{-10}$	90	0	YOCK	75	ELEC $\pm$	$Q > 7e \text{ or } < -7e$
$> 6. \times 10^{-9}$		5	142 YOCK	74	CNTR	$m > 6 \text{ GeV}$
$< 3.0 \times 10^{-8}$		0	DARDO	72	CNTR	
$< 1.5 \times 10^{-9}$		0	TONWAR	72	CNTR	$m > 10 \text{ GeV}$
$< 3.0 \times 10^{-10}$		0	BJORNBOE	68	CNTR	$m > 5 \text{ GeV}$
$< 5.0 \times 10^{-11}$	90	0	JONES	67	ELEC	$m = 5-15 \text{ GeV}$

134 SAITO 90 candidates carry about 450 MeV/nucleon. Cannot be accounted for by conventional backgrounds. Consistent with strange quark matter hypothesis.

135 MINCER 85 is high statistics study of calorimeter signals delayed by 20–200 ns. Calibration with AGS beam shows they can be accounted for by rare fluctuations in signals from low-energy hadrons in the shower. Claim that previous delayed signals including BJORNBOE 68, DARDO 72, BHAT 82, SAKUYAMA 83B below may be due to this fake effect.

136 SAKUYAMA 83B analyzed 6000 extended air shower events. Increase of delayed particles and change of lateral distribution above  $10^{17}$  eV may indicate production of very heavy parent at top of atmosphere.

- 137 BHAT 82 observed 12 events with delay  $> 2 \times 10^{-8}$  s and with more than 40 particles. 1 eV has good hadron shower. However all events are delayed in only one of two detectors in cloud chamber, and could not be due to strongly interacting massive particle.
- 138 MARINI 82 applied PEP-counter for TOF. Above limit is for velocity = 0.54 of light. Limit is inconsistent with YOCK 80 YOCK 81 events if isotropic dependence on zenith angle is assumed.
- 139 YOCK 81 saw another 3 events with  $Q = \pm 1$  and  $m$  about  $4.5m_p$  as well as 2 events with  $m > 5.3m_p$ ,  $Q = \pm 0.75 \pm 0.05$  and  $m > 2.8m_p$ ,  $Q = \pm 0.70 \pm 0.05$  and 1 event with  $m = (9.3 \pm 3.)m_p$ ,  $Q = \pm 0.89 \pm 0.06$  as possible heavy candidates.
- 140 YOCK 80 events are with charge exactly or approximately equal to unity.
- 141 BHAT 78 is at Kolar gold fields. Limit is for  $\tau > 10^{-6}$  s.
- 142 YOCK 74 events could be tritons.

## Superheavy Particle (Quark Matter) Flux in Cosmic Rays

<u>VALUE</u> ( $\text{cm}^{-2}\text{sr}^{-1}\text{s}^{-1}$ )	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>					
$<5 \times 10^{-16}$	90		143 AMBROSIO	00B MCRO	$m > 5 \times 10^{14}$ GeV
$<1.8 \times 10^{-12}$	90		144 ASTONE	93 CNTR	$m \geq 1.5 \times 10^{-13}$ gram
$<1.1 \times 10^{-14}$	90		145 AHLEN	92 MCRO	$10^{-10} < m < 0.1$ gram
$<2.2 \times 10^{-14}$	90	0	146 NAKAMURA	91 PLAS	$m > 10^{11}$ GeV
$<6.4 \times 10^{-16}$	90	0	147 ORITO	91 PLAS	$m > 10^{12}$ GeV
$<2.0 \times 10^{-11}$	90		148 LIU	88 BOLO	$m > 1.5 \times 10^{-13}$ gram
$<4.7 \times 10^{-12}$	90		149 BARISH	87 CNTR	$1.4 \times 10^8 < m < 10^{12}$ GeV
$<3.2 \times 10^{-11}$	90	0	150 NAKAMURA	85 CNTR	$m > 1.5 \times 10^{-13}$ gram
$<3.5 \times 10^{-11}$	90	0	151 ULLMAN	81 CNTR	Planck-mass $10^{19}$ GeV
$<7. \times 10^{-11}$	90	0	151 ULLMAN	81 CNTR	$m \leq 10^{16}$ GeV

- 143 AMBROSIO 00B searched for quark matter ("nuclearites") in the velocity range  $(10^{-5}-1)c$ . The listed limit is for  $2 \times 10^{-3}c$ .
- 144 ASTONE 93 searched for quark matter ("nuclearites") in the velocity range  $(10^{-3}-1)c$ . Their Table 1 gives a compilation of searches for nuclearites.
- 145 AHLEN 92 searched for quark matter ("nuclearites"). The bound applies to velocity  $< 2.5 \times 10^{-3}c$ . See their Fig. 3 for other velocity/c and heavier mass range.
- 146 NAKAMURA 91 searched for quark matter in the velocity range  $(4 \times 10^{-5}-1)c$ .
- 147 ORITO 91 searched for quark matter. The limit is for the velocity range  $(10^{-4}-10^{-3})c$ .
- 148 LIU 88 searched for quark matter ("nuclearites") in the velocity range  $(2.5 \times 10^{-3}-1)c$ . A less stringent limit of  $5.8 \times 10^{-11}$  applies for  $(1-2.5) \times 10^{-3}c$ .
- 149 BARISH 87 searched for quark matter ("nuclearites") in the velocity range  $(2.7 \times 10^{-4}-5 \times 10^{-3})c$ .
- 150 NAKAMURA 85 at KEK searched for quark-matter. These might be lumps of strange quark matter with roughly equal numbers of  $u$ ,  $d$ ,  $s$  quarks. These lumps or nuclearites were assumed to have velocity of  $(10^{-4}-10^{-3})c$ .
- 151 ULLMAN 81 is sensitive for heavy slow singly charge particle reaching earth with vertical velocity 100–350 km/s.

## Highly Ionizing Particle Flux

<u>VALUE</u> ( $\text{m}^{-2}\text{yr}^{-1}$ )	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>					
<0.4	95	0	KINOSHITA	81B PLAS	$Z/\beta$ 30–100

## REFERENCES FOR Searches for WIMPs and Other Particles

AKERIB	06	PR D73 011102R	D.S. Akerib <i>et al.</i>	(CDMS Collab.)
SHIMIZU	06A	PL B633 195	Y. Shimizu <i>et al.</i>	
AKERIB	05	PR D72 052009	D.S. Akerib <i>et al.</i>	(CDMS Collab.)
ALNER	05	PL B616 17	G.J. Alner <i>et al.</i>	(UK Dark Matter Collab.)
BARNABE-HE...	05	PL B624 186	M. Barnabe-Heider <i>et al.</i>	(PICASSO Collab.)
BENOIT	05	PL B616 25	A. Benoit <i>et al.</i>	(EDELWEISS Collab.)
GIRARD	05	PL B621 233	T.A. Girard <i>et al.</i>	(SIMPLE Collab.)
GULIANI	05	PRL 95 101301	F. Giuliani	
GULIANI	05A	PR D71 123503	F. Giuliani, T.A. Girard	
KLAPDOR-K...	05	PL B609 226	H.V. Klapdor-Kleingrothaus, I.V. Krivosheina, C. Tomei	
AKTAS	04C	EPJ C36 413	A. Atkas <i>et al.</i>	(H1 Collab.)
GULIANI	04	PL B588 151	F. Giuliani, T.A. Girard	
GULIANI	04A	PRL 93 161301	F. Giuliani	
MIUCHI	03	ASP 19 135	K. Miuchi <i>et al.</i>	
TAKEDA	03	PL B572 145	A. Takeda <i>et al.</i>	
ANGLOHER	02	ASP 18 43	G. Angloher <i>et al.</i>	(CRESST Collab.)
BELLI	02	PR D66 043503	P. Belli <i>et al.</i>	
BERNABEI	02C	EPJ C23 61	R. Bernabei <i>et al.</i>	(DAMA Collab.)
GREEN	02	PR D66 083003	A.M. Green	
JAVORSEK	02	PR D65 072003	D. Javorsek II <i>et al.</i>	
JAVORSEK	01	PR D64 012005	D. Javorsek II <i>et al.</i>	
JAVORSEK	01B	PRL 87 231804	D. Javorsek II <i>et al.</i>	
SMITH	01	PR D64 043502	D. Smith, N. Weiner	
ULLIO	01	JHEP 0107 044	P. Ullio, M. Kamionkowski, P. Vogel	
ABBIENDI	00D	EPJ C13 197	G. Abbiendi <i>et al.</i>	(OPAL Collab.)
AMBROSIO	00B	EPJ C13 453	M. Ambrosio <i>et al.</i>	(MACRO Collab.)
BENOIT	00	PL B479 8	A. Benoit <i>et al.</i>	(EDELWEISS Collab.)
BERNABEI	00D	NJP 2 15	R. Bernabei <i>et al.</i>	(DAMA Collab.)
COLLAR	00	PRL 85 3083	J.I. Collar <i>et al.</i>	(SIMPLE Collab.)
ABE	99F	PRL 82 2038	F. Abe <i>et al.</i>	(CDF Collab.)
AMBROSIO	99	PR D60 082002	M. Ambrosio <i>et al.</i>	(Macro Collab.)
BERNABEI	99	PL B450 448	R. Bernabei <i>et al.</i>	(DAMA Collab.)
BERNABEI	99D	PRL 83 4918	R. Bernabei <i>et al.</i>	(DAMA Collab.)
BRHLIK	99	PL B464 303	M. Brhlik, L. Roszkowski	
DERBIN	99	PAN 62 1886	A.V. Derbin <i>et al.</i>	
		Translated from YAF 62 2034.		
ACKERSTAFF	98P	PL B433 195	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
KLIMENTKO	98	JETPL 67 875	A.A. Klimenko <i>et al.</i>	
		Translated from ZETFP 67 835.		
ABE	97G	PR D55 R5263	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	97D	PL B396 315	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACKERSTAFF	97B	PL B391 210	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
ADAMS	97B	PRL 79 4083	J. Adams <i>et al.</i>	(FNAL KTeV Collab.)
BARATE	97K	PL B405 379	R. Barate <i>et al.</i>	(ALEPH Collab.)
SARSA	97	PR D56 1856	M.L. Sarsa <i>et al.</i>	(ZARA)
ALESSAND...	96	PL B384 316	A. Alessandrello <i>et al.</i>	(MILA, MILAI, SASSO)
BELLI	96	PL B387 222	P. Belli <i>et al.</i>	(DAMA Collab.)
Also		PL B389 783 (erratum)	P. Belli <i>et al.</i>	(DAMA Collab.)
BELLI	96C	NC 19C 537	P. Belli <i>et al.</i>	(DAMA Collab.)
BERNABEI	96	PL B389 757	R. Bernabei <i>et al.</i>	(DAMA Collab.)
COLLAR	96	PRL 76 331	J.I. Collar	(SCUC)
SARSA	96	PL B386 458	M.L. Sarsa <i>et al.</i>	(ZARA)
Also		PR D56 1856	M.L. Sarsa <i>et al.</i>	(ZARA)
SMITH	96	PL B379 299	P.F. Smith <i>et al.</i>	(RAL, SHEF, LOIC+)
SNOWDEN-...	96	PRL 76 332	D.P. Snowden-Ifft, E.S. Freeman, P.B. Price	(UCB)
AKERS	95R	ZPHY C67 203	R. Akers <i>et al.</i>	(OPAL Collab.)
GALLAS	95	PR D52 6	E. Gallas <i>et al.</i>	(MSU, FNAL, MIT, FLOR)
GARCIA	95	PR D51 1458	E. Garcia <i>et al.</i>	(ZARA, SCUC, PNL)
QUEENBY	95	PL B351 70	J.J. Queenby <i>et al.</i>	(LOIC, RAL, SHEF+)
SNOWDEN-...	95	PRL 74 4133	D.P. Snowden-Ifft, E.S. Freeman, P.B. Price	(UCB)
Also		PL 76 331	J.I. Collar	(SCUC)
Also		PRL 76 332	D.P. Snowden-Ifft, E.S. Freeman, P.B. Price	(UCB)
BECK	94	PL B336 141	M. Beck <i>et al.</i>	(MPIH, KIAE, SASSO)
RAM	94	PR D49 3120	S. Ram <i>et al.</i>	(TELA, TRIU)
ABE	93G	PRL 71 2542	F. Abe <i>et al.</i>	(CDF Collab.)
ASTONE	93	PR D47 4770	P. Astone <i>et al.</i>	(ROMA, ROMAI, CATA, FRAS)
BUSKULIC	93C	PL B303 198	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
YAMAGATA	93	PR D47 1231	T. Yamagata, Y. Takamori, H. Utsunomiya	(KONAN)
ABE	92J	PR D46 R1889	F. Abe <i>et al.</i>	(CDF Collab.)

AHLEN	92	PRL 69 1860	S.P. Ahlen <i>et al.</i>	(MACRO Collab.)
BACCI	92	PL B293 460	C. Bacci <i>et al.</i>	(Beijing-Roma-Saclay Collab.)
VERKERK	92	PRL 68 1116	P. Verkerk <i>et al.</i>	(ENSP, SACL, PAST)
AKESSON	91	ZPHY C52 219	T. Akesson <i>et al.</i>	(HELIOS Collab.)
NAKAMURA	91	PL B263 529	S. Nakamura <i>et al.</i>	
ORITO	91	PRL 66 1951	S. Orito <i>et al.</i>	(ICEPP, WASCR, NIHO, ICRR)
REUSSER	91	PL B255 143	D. Reusser <i>et al.</i>	(NEUC, CIT, PSI)
ADACHI	90C	PL B244 352	I. Adachi <i>et al.</i>	(TOPAZ Collab.)
ADACHI	90E	PL B249 336	I. Adachi <i>et al.</i>	(TOPAZ Collab.)
AKRAWY	90O	PL B252 290	M.Z. Akrawy <i>et al.</i>	(OPAL Collab.)
HEMMICK	90	PR D41 2074	T.K. Hemmick <i>et al.</i>	(ROCH, MICH, OHIO+)
SAITO	90	PRL 65 2094	T. Saito <i>et al.</i>	(ICRR, KOBE)
NAKAMURA	89	PR D39 1261	T.T. Nakamura <i>et al.</i>	(KYOT, TMT)
NORMAN	89	PR D39 2499	E.B. Norman <i>et al.</i>	(LBL)
BERNSTEIN	88	PR D37 3103	R.M. Bernstein <i>et al.</i>	(STAN, WISC)
CALDWELL	88	PRL 61 510	D.O. Caldwell <i>et al.</i>	(UCSB, UCB, LBL)
LIU	88	PRL 61 271	G. Liu, B. Barish	
BARISH	87	PR D36 2641	B.C. Barish, G. Liu, C. Lane	(CIT)
NORMAN	87	PRL 58 1403	E.B. Norman, S.B. Gazes, D.A. Bennett	(LBL)
BADIER	86	ZPHY C31 21	J. Badier <i>et al.</i>	(NA3 Collab.)
MINCER	85	PR D32 541	A. Mincer <i>et al.</i>	(UMD, GMAS, NSF)
NAKAMURA	85	PL 161B 417	K. Nakamura <i>et al.</i>	(KEK, INUS)
THON	85	PR D31 451	J.L. Thron <i>et al.</i>	(YALE, FNAL, IOWA)
SAKUYAMA	83B	LNC 37 17	H. Sakuyama, N. Suzuki	(MEIS)
Also		LNC 36 389	H. Sakuyama, K. Watanabe	(MEIS)
Also		NC 78A 147	H. Sakuyama, K. Watanabe	(MEIS)
Also		NC 6C 371	H. Sakuyama, K. Watanabe	(MEIS)
BHAT	82	PR D25 2820	P.N. Bhat <i>et al.</i>	(TATA)
KINOSHITA	82	PRL 48 77	K. Kinoshita, P.B. Price, D. Fryberger	(UCB+)
MARINI	82	PR D26 1777	A. Marini <i>et al.</i>	(FRAS, LBL, NWES, STAN+)
SMITH	82B	NP B206 333	P.F. Smith <i>et al.</i>	(RAL)
KINOSHITA	81B	PR D24 1707	K. Kinoshita, P.B. Price	(UCB)
LOSECCO	81	PL 102B 209	J.M. LoSecco <i>et al.</i>	(MICH, PENN, BNL)
ULLMAN	81	PRL 47 289	J.D. Ullman	(LEHM, BNL)
YOCK	81	PR D23 1207	P.C.M. Yock	(AUCK)
BARTEL	80	ZPHY C6 295	W. Bartel <i>et al.</i>	(JADE Collab.)
BUSSIÈRE	80	NP B174 1	A. Bussiere <i>et al.</i>	(BGNA, SACL, LAPP)
YOCK	80	PR D22 61	P.C.M. Yock	(AUCK)
ARMITAGE	79	NP B150 87	J.C.M. Armitage <i>et al.</i>	(CERN, DARE, FOM+)
BOZZOLI	79	NP B159 363	W. Bozzoli <i>et al.</i>	(BGNA, LAPP, SACL+)
GOODMAN	79	PR D19 2572	J.A. Goodman <i>et al.</i>	(UMD)
SMITH	79	NP B149 525	P.F. Smith, J.R.J. Bennett	(RHEL)
BHAT	78	PRAM 10 115	P.N. Bhat, P.V. Ramana Murthy	(TATA)
CARROLL	78	PRL 41 777	A.S. Carroll <i>et al.</i>	(BNL, PRIN)
CUTTS	78	PRL 41 363	D. Cutts <i>et al.</i>	(BROW, FNAL, ILL, BARI+)
VIDAL	78	PL 77B 344	R.A. Vidal <i>et al.</i>	(COLU, FNAL, STON+)
ALEKSEEV	76	SJNP 22 531	G.D. Alekseev <i>et al.</i>	(JINR)
ALEKSEEV	76	Translated from YAF 22 1021.	G.D. Alekseev <i>et al.</i>	
ALEKSEEV	76B	SJNP 23 633	Translated from YAF 23 1190.	(JINR)
BALDIN	76	SJNP 22 264	B.Y. Baldin <i>et al.</i>	(JINR)
BALDIN	76	Translated from YAF 22 512.		
BRIATORE	76	NC 31A 553	L. Briatore <i>et al.</i>	(LCGT, FRAS, FREIB)
GUSTAFSON	76	PRL 37 474	H.R. Gustafson <i>et al.</i>	(MICH)
ALBROW	75	NP B97 189	M.G. Albrow <i>et al.</i>	(CERN, DARE, FOM+)
FRANKEL	75	PR D12 2561	S. Frankel <i>et al.</i>	(PENN, FNAL)
JOVANOV...	75	PL 56B 105	J.V. Jovanovich <i>et al.</i>	(MANI, AACH, CERN+)
YOCK	75	NP B86 216	P.C.M. Yock	(AUCK, SLAC)
APPEL	74	PRL 32 428	J.A. Appel <i>et al.</i>	(COLU, FNAL)
FRANKEL	74	PR D9 1932	S. Frankel <i>et al.</i>	(PENN, FNAL)
YOCK	74	NP B76 175	P.C.M. Yock	(AUCK)
ALPER	73	PL 46B 265	B. Alper <i>et al.</i>	(CERN, LIVP, LUND, BOHR+)
LEIPUNER	73	PRL 31 1226	L.B. Leipuner <i>et al.</i>	(BNL, YALE)
DARDO	72	NC 9A 319	M. Dardo <i>et al.</i>	(TORI)
TONWAR	72	JPA 5 569	S.C. Tonwar, S. Naranan, B.V. Sreekantan	(TATA)
ANTIROV	71B	NP B31 235	Y.M. Antipov <i>et al.</i>	(SERP)
ANTIROV	71C	PL 34B 164	Y.M. Antipov <i>et al.</i>	(SERP)
BINON	69	PL 30B 510	F.G. Binon <i>et al.</i>	(SERP)
BJORNBOE	68	NC B53 241	J. Bjornboe <i>et al.</i>	(BOHR, TATA, BERN+)
JONES	67	PR 164 1584	L.W. Jones	(MICH, WISC, LBL, UCLA, MINN+)
DORFAN	65	PRL 14 999	D.E. Dorfan <i>et al.</i>	(COLU)